

1. Let $P = P(t)$ be the population at time t , and P_0 the initial population. From $dP/dt = kP$ we obtain $P = P_0e^{kt}$. Using $P(5) = 2P_0$ we find $k = \frac{1}{5} \ln 2$ and $P = P_0e^{(\ln 2)t/5}$. Setting $P(t) = 3P_0$ we have $3 = e^{(\ln 2)t/5}$, so

$$\ln 3 = \frac{(\ln 2)t}{5} \quad \text{and} \quad t = \frac{5 \ln 3}{\ln 2} \approx 7.9 \text{ years.}$$

Setting $P(t) = 4P_0$ we have $4 = e^{(\ln 2)t/5}$, so

$$\ln 4 = \frac{(\ln 2)t}{5} \quad \text{and} \quad t \approx 10 \text{ years.}$$

3. Let $P = P(t)$ be the population at time t . Then $dP/dt = kP$ and $P = ce^{kt}$. From $P(0) = c = 500$ we see that $P = 500e^{kt}$. Since 15% of 500 is 75, we have $P(10) = 500e^{10k} = 575$. Solving for k , we get $k = \frac{1}{10} \ln \frac{575}{500} = \frac{1}{10} \ln 1.15$. When $t = 30$,

$$P(30) = 500e^{(1/10)(\ln 1.15)30} = 500e^{3 \ln 1.15} = 760 \text{ years}$$

and

$$P'(30) = kP(30) = \frac{1}{10}(\ln 1.15)760 = 10.62 \text{ persons/year.}$$

5. Let $A = A(t)$ be the amount of lead present at time t . From $dA/dt = kA$ and $A(0) = 1$ we obtain $A = e^{kt}$. Using $A(3.3) = 1/2$ we find $k = \frac{1}{3.3} \ln(1/2)$. When 90% of the lead has decayed, 0.1 grams will remain. Setting $A(t) = 0.1$ we have $e^{t(1/3.3) \ln(1/2)} = 0.1$, so

$$\frac{t}{3.3} \ln \frac{1}{2} = \ln 0.1 \quad \text{and} \quad t = \frac{3.3 \ln 0.1}{\ln(1/2)} \approx 10.96 \text{ hours.}$$

7. Setting $A(t) = 50$ in Problem 6 we obtain $50 = 100e^{kt}$, so

$$kt = \ln \frac{1}{2} \quad \text{and} \quad t = \frac{\ln(1/2)}{(1/6) \ln 0.97} \approx 136.5 \text{ hours.}$$

9. Let $I = I(t)$ be the intensity, t the thickness, and $I(0) = I_0$. If $dI/dt = kI$ and $I(3) = 0.25I_0$, then $I = I_0e^{kt}$, $k = \frac{1}{3} \ln 0.25$, and $I(15) = 0.00098I_0$.

11. Assume that $A = A_0e^{kt}$ and $k = -0.00012378$. If $A(t) = 0.145A_0$ then $t \approx 15,600$ years.

13. Assume that $dT/dt = k(T - 10)$ so that $T = 10 + ce^{kt}$. If $T(0) = 70^\circ$ and $T(1/2) = 50^\circ$ then $c = 60$ and $k = 2 \ln(2/3)$ so that $T(1) = 36.67^\circ$. If $T(t) = 15^\circ$ then $t = 3.06$ minutes.

15. We use the fact that the boiling temperature for water is 100° C. Now assume that $dT/dt = k(T - 100)$ so that $T = 100 + ce^{kt}$. If $T(0) = 20^\circ$ and $T(1) = 22^\circ$, then $c = -80$ and $k = \ln(39/40) \approx -0.0253$. Then $T(t) = 100 - 80e^{-0.0253t}$, and when $T = 90$, $t = 82.1$ seconds. If $T(t) = 98^\circ$ then $t = 145.7$ seconds.

17. Using separation of variables to solve $dT/dt = k(T - T_m)$ we get $T(t) = T_m + ce^{kt}$. Using $T(0) = 70$ we find $c = 70 - T_m$, so $T(t) = T_m + (70 - T_m)e^{kt}$. Using the given observations, we obtain

$$T\left(\frac{1}{2}\right) = T_m + (70 - T_m)e^{k/2} = 110$$

$$T(1) = T_m + (70 - T_m)e^k = 145.$$

Then, from the first equation, $e^{k/2} = (110 - T_m)/(70 - T_m)$ and

$$e^k = (e^{k/2})^2 = \left(\frac{110 - T_m}{70 - T_m}\right)^2 = \frac{145 - T_m}{70 - T_m}$$

$$\frac{(110 - T_m)^2}{70 - T_m} = 145 - T_m$$

$$12100 - 220T_m + T_m^2 = 10150 - 215T_m + T_m^2$$

$$T_m = 390.$$

The temperature in the oven is 390° .

19. Identifying $T_m = 70$, the differential equation is $dT/dt = k(T - 70)$. Assuming $T(0) = 98.6$ and separating variables we find $T(t) = 70 + 28.9e^{kt}$. If $t_1 > 0$ is the time of discovery of the body, then

$$T(t_1) = 70 + 28.6e^{kt_1} = 85 \quad \text{and} \quad T(t_1 + 1) = 70 + 28.6e^{k(t_1+1)} = 80.$$

Therefore $e^{kt_1} = 15/28.6$ and $e^{k(t_1+1)} = 10/28.6$. This implies

$$e^k = \frac{10}{28.6} e^{-kt_1} = \frac{10}{28.6} \cdot \frac{28.6}{15} = \frac{2}{3},$$

so $k = \ln \frac{2}{3} \approx -0.405465108$. Therefore

$$t_1 = \frac{1}{k} \ln \frac{15}{28.6} \approx 1.5916 \approx 1.6.$$

Death took place about 1.6 hours prior to the discovery of the body.

21. From $dA/dt = 4 - A/50$ we obtain $A = 200 + ce^{-t/50}$. If $A(0) = 30$ then $c = -170$ and $A = 200 - 170e^{-t/50}$.
23. From $dA/dt = 10 - A/100$ we obtain $A = 1000 + ce^{-t/100}$. If $A(0) = 0$ then $c = -1000$ and $A(t) = 1000 - 1000e^{-t/100}$.

25. From

$$\frac{dA}{dt} = 10 - \frac{10A}{500 - (10 - 5)t} = 10 - \frac{2A}{100 - t}$$

we obtain $A = 1000 - 10t + c(100 - t)^2$. If $A(0) = 0$ then $c = -\frac{1}{10}$. The tank is empty in 100 minutes.

27. From

$$\frac{dA}{dt} = 3 - \frac{4A}{100 + (6 - 4)t} = 3 - \frac{2A}{50 + t}$$

we obtain $A = 50 + t + c(50 + t)^{-2}$. If $A(0) = 10$ then $c = -100,000$ and $A(30) = 64.38$ pounds.

29. Assume $L di/dt + Ri = E(t)$, $L = 0.1$, $R = 50$, and $E(t) = 50$ so that $i = \frac{3}{5} + ce^{-500t}$. If $i(0) = 0$ then $c = -3/5$ and $\lim_{t \rightarrow \infty} i(t) = 3/5$.

31. Assume $R dq/dt + (1/C)q = E(t)$, $R = 200$, $C = 10^{-4}$, and $E(t) = 100$ so that $q = 1/100 + ce^{-50t}$. If $q(0) = 0$ then $c = -1/100$ and $i = \frac{1}{2}e^{-50t}$.

33. For $0 \leq t \leq 20$ the differential equation is $20 di/dt + 2i = 120$. An integrating factor is $e^{t/10}$, so $(d/dt)[e^{t/10}i] = 6e^{t/10}$ and $i = 60 + c_1e^{-t/10}$. If $i(0) = 0$ then $c_1 = -60$ and $i = 60 - 60e^{-t/10}$. For $t > 20$ the differential equation is $20 di/dt + 2i = 0$ and $i = c_2e^{-t/10}$. At $t = 20$ we want $c_2e^{-2} = 60 - 60e^{-2}$ so that $c_2 = 60(e^2 - 1)$. Thus

$$i(t) = \begin{cases} 60 - 60e^{-t/10}, & 0 \leq t \leq 20 \\ 60(e^2 - 1)e^{-t/10}, & t > 20. \end{cases}$$

35. (a) From $m dv/dt = mg - kv$ we obtain $v = mg/k + ce^{-kt/m}$. If $v(0) = v_0$ then $c = v_0 - mg/k$ and the solution of the initial-value problem is

$$v(t) = \frac{mg}{k} + \left(v_0 - \frac{mg}{k}\right) e^{-kt/m}.$$

(b) As $t \rightarrow \infty$ the limiting velocity is mg/k .

(c) From $ds/dt = v$ and $s(0) = 0$ we obtain

$$s(t) = \frac{mg}{k}t - \frac{m}{k} \left(v_0 - \frac{mg}{k}\right) e^{-kt/m} + \frac{m}{k} \left(v_0 - \frac{mg}{k}\right).$$

39. (a) The differential equation is first-order and linear. Letting $b = k/\rho$, the integrating factor is $e^{\int 3b dt/(bt+r_0)} = (r_0 + bt)^3$. Then

$$\frac{d}{dt}[(r_0 + bt)^3 v] = g(r_0 + bt)^3 \quad \text{and} \quad (r_0 + bt)^3 v = \frac{g}{4b}(r_0 + bt)^4 + c.$$


The solution of the differential equation is $v(t) = (g/4b)(r_0 + bt) + c(r_0 + bt)^{-3}$. Using $v(0) = 0$ we find $c = -gr_0^4/4b$, so that

$$v(t) = \frac{g}{4b}(r_0 + bt) - \frac{gr_0^4}{4b(r_0 + bt)^3} = \frac{g\rho}{4k}\left(r_0 + \frac{k}{\rho}t\right) - \frac{g\rho r_0^4}{4k(r_0 + kt/\rho)^3}.$$

- (b) Integrating $dr/dt = k/\rho$ we get $r = kt/\rho + c$. Using $r(0) = r_0$ we have $c = r_0$, so $r(t) = kt/\rho + r_0$.
 (c) If $r = 0.007$ ft when $t = 10$ s, then solving $r(10) = 0.007$ for k/ρ , we obtain $k/\rho = -0.0003$ and $r(t) = 0.01 - 0.0003t$. Solving $r(t) = 0$ we get $t = 33.3$, so the raindrop will have evaporated completely at 33.3 seconds.

41. (a) From $dP/dt = (k_1 - k_2)P$ we obtain $P = P_0 e^{(k_1 - k_2)t}$ where $P_0 = P(0)$.
 (b) If $k_1 > k_2$ then $P \rightarrow \infty$ as $t \rightarrow \infty$. If $k_1 = k_2$ then $P = P_0$ for every t . If $k_1 < k_2$ then $P \rightarrow 0$ as $t \rightarrow \infty$.

43. (a) Solving $r - kx = 0$ for x we find the equilibrium solution $x = r/k$. When $x < r/k$, $dx/dt > 0$ and when $x > r/k$, $dx/dt < 0$. From the phase portrait we see that $\lim_{t \rightarrow \infty} x(t) = r/k$.



- (b) From $dx/dt = r - kx$ and $x(0) = 0$ we obtain $x = r/k - (r/k)e^{-kt}$ so that $x \rightarrow r/k$ as $t \rightarrow \infty$. If $x(T) = r/2k$ then $T = (\ln 2)/k$.

