

Volume

p. 528: 9-25 odd, 39, 41-44



9. A cross-section is a disk with radius  $x+1$ , so its area is

$$A(x) = \pi(x+1)^2 = \pi(x^2 + 2x + 1).$$

$$= \pi(x+1)^2 = \pi(x^2 + 2x + 1).$$

$$V = \int_0^2 A(x) dx = \int_0^2 \pi(x^2 + 2x + 1) dx$$

$$= \pi \left[ \frac{1}{3}x^3 + x^2 + x \right]_0^2 = \pi \left( \frac{8}{3} + 4 + 2 \right) = \frac{26}{3} \pi$$





11. A cross-section is a disk with radius  $\sqrt{x-1}$ , so its area is

$$A(x) = \pi(\sqrt{x-1})^2 = \pi(x-1).$$

$$V = \int_1^5 A(x) dx = \int_1^5 \pi(x-1) dx$$

$$= \pi \left[ \frac{1}{2}x^2 - x \right]_1^5$$

$$= \pi \left[ \left( \frac{25}{2} - 5 \right) - \left( \frac{1}{2} - 1 \right) \right] = 8\pi$$





13. A cross-section is a disk with radius  $2\sqrt{y}$  so its area is

$$A(y) = \pi(2\sqrt{y})^2.$$

$$V = \int_0^9 A(y) dy = \pi \int_0^9 (2\sqrt{y})^2 dy = 4\pi \int_0^9 y dy$$

$$= 4\pi \left[ \frac{1}{2}y^2 \right]_0^9 = 2\pi(81) = 162\pi$$





15. A cross-section is a washer (annulus) with inner radius  $x^3$  and outer radius  $x$  so its area is

$$A(x) = \pi(x)^2 - \pi(x^3)^2 = \pi(x^2 - x^6).$$

$$V = \int_0^1 A(x) dx = \pi \int_0^1 (x^2 - x^6) dx$$

$$= \pi \left[ \frac{1}{3}x^3 - \frac{1}{7}x^7 \right]_0^1 = \pi \left( \frac{1}{3} - \frac{1}{7} \right) = \frac{4}{21} \pi$$





17. A cross-section is a washer with inner radius  $y^2$  and outer radius  $2y$  so its area is

$$A(y) = \pi(2y)^2 - \pi(y^2)^2 = \pi(4y^2 - y^4).$$

$$V = \int_0^2 A(y) dy = \pi \int_0^2 (4y^2 - y^4) dy$$


$$= \pi \left[ \frac{4}{3}y^3 - \frac{1}{5}y^5 \right]_0^2 = \pi \left( \frac{32}{3} - \frac{32}{5} \right) = \frac{64}{15} \pi$$



19. A cross-section is a washer with inner radius  $1 - \sqrt{x}$  and outer radius  $1 - x^2$  so its area is


$$\begin{aligned} A(x) &= \pi \left[ (1 - x^2)^2 - (1 - \sqrt{x})^2 \right] \\ &= \pi \left[ (1 - 2x^2 + x^4) - (1 - 2\sqrt{x} + x) \right] \\ &= \pi (x^4 - 2x^2 + 2\sqrt{x} - x). \end{aligned}$$

$$V = \int_0^1 A(y) dy = \pi \int_0^1 (x^4 - 2x^2 + 2x^{1/2} - x) dx = \pi \left[ \frac{1}{5}x^5 - \frac{2}{3}x^3 + \frac{4}{3}x^{3/2} - \frac{1}{2}x^2 \right]_0^1 = \pi \left( \frac{1}{5} - \frac{2}{3} + \frac{4}{3} - \frac{1}{2} \right) = \frac{11}{30} \pi$$



21. A cross-section is a washer with inner radius  $(1 + \sec x) - 1 = \sec x$  and outer radius  $3 - 1 = 2$ , so its area is  $A(x) = \pi [2^2 - \sec^2 x] = \pi(4 - \sec^2 x)$ .


$$\begin{aligned} V &= \int_{-\pi/3}^{\pi/3} A(x) dx = \pi \int_{-\pi/3}^{\pi/3} (4 - \sec^2 x) dx \\ &= 2\pi \int_0^{\pi/3} (4 - \sec^2 x) dx \quad [\text{by symmetry}] \\ &= 2\pi [4x - \tan x]_0^{\pi/3} = 2\pi \left[ \left( \frac{4\pi}{3} - \sqrt{3} \right) - 0 \right] \\ &= 2\pi \left( \frac{4\pi}{3} - \sqrt{3} \right) \end{aligned}$$



23. A cross-section is a washer with inner radius  $2 - 1$  and outer radius  $2 - \sqrt[3]{y}$ , so its area is

$$A(y) = \pi \left[ (2 - \sqrt[3]{y})^2 - (2 - 1)^2 \right] = \pi \left[ 4 - 4\sqrt[3]{y} + \sqrt[3]{y^2} - 1 \right].$$


$$V = \int_0^1 A(y) dy = \pi \int_0^1 (3 - 4y^{1/3} + y^{2/3}) dy = \pi \left[ 3y - 3y^{4/3} + \frac{3}{5}y^{5/3} \right]_0^1 = \pi \left( 3 - 3 + \frac{3}{5} \right) = \frac{3}{5} \pi$$





25. From the symmetry of the curves, we see that they intersect at  $x = \frac{1}{2}$  and so  $y^2 = \frac{1}{2} \Leftrightarrow y = \pm\sqrt{\frac{1}{2}}$ . A cross-section is a washer with inner radius  $3 - (1 - y^2)$  and outer radius  $3 - y^2$ , so its area is

$$\begin{aligned} A(y) &= \pi \left[ (3 - y^2)^2 - (2 + y^2)^2 \right] \\ &= \pi \left[ (9 - 6y^2 + y^4) - (4 + 4y^2 + y^4) \right] = \pi(5 - 10y^2). \end{aligned}$$

$$\begin{aligned} V &= \int_{-\sqrt{1/2}}^{\sqrt{1/2}} A(y) dy = 2 \int_0^{\sqrt{1/2}} 5\pi(1 - 2y^2) dy \quad [\text{by symmetry}] \\ &= 10\pi \left[ y - \frac{2}{3}y^3 \right]_0^{\sqrt{1/2}} = 10\pi \left( \frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{6} \right) = 10\pi \left( \frac{\sqrt{2}}{3} \right) = \frac{10}{3} \sqrt{2} \pi \end{aligned}$$



39. The area of this region is  $A(y) = \pi(2 - y^{1/3})^2$ , so the volume is  $V = \int_0^8 \pi(2 - y^{1/3})^2 dy$ , option (B).



41. A cross-section is a washer with inner radius  $x-1$  and outer radius  $2 \ln x$ , so the area is

$A(x) = \pi[(2 \ln x)^2 - (x-1)^2] = \pi(4(\ln x)^2 - x^2 + 2x - 1)$ . The curves intersect at  $x = 1$  and  $x = a \approx 3.512862417$ . Therefore,

$$V = \pi \int_1^a A(x) dx = \pi \int_1^a (4(\ln x)^2 - x^2 + 2x - 1) dx \stackrel{\text{CAS}}{\approx} 5.298, \text{ which is option (A).}$$

42. A cross-section is a washer with inner radius  $y = \sqrt{x} \Rightarrow x = y^2$  and outer radius

$y = 6 - x \Rightarrow x = 6 - y$ , so its area is  $A(y) = \pi(6 - y)^2 - \pi(y^2)^2$ . The volume of the resulting solid is

$$\text{(A) } \pi \int_0^2 [(6 - y)^2 - (y^2)^2] dy.$$

43. A cross-section is a washer with radius  $\sqrt{x}$  so the area is  $A(x) = \pi(\sqrt{x})^2 = \pi x$ , and the volume is

$$V = \pi \int_0^4 A(x) dx = \pi \int_0^4 x dx = \pi \left[ \frac{1}{2} x^2 \right]_0^4 = \pi \left( \frac{1}{2} (16) - 0 \right) = 8\pi, \text{ which is option (D).}$$

44. A cross-section is a washer with radius  $\sec x$  so the area is  $A(x) = \pi \sec^2 x$  and the volume is

$$V = \pi \int_0^{\pi/4} A(x) dx = \pi \int_0^{\pi/4} \sec^2 x dx = \pi [\tan x]_0^{\pi/4} = \pi \left[ \tan \frac{\pi}{4} - \tan 0 \right] = \pi(1 - 0) = \pi, \text{ option (A).}$$